Differences in the Composition of Soils Under Open and Canopy Conditions at Two Sites Close-in to the Great Canadian Oil Sands Operation, Fort McMurray, Alberta

Project LS 3.4.1
April 1980

FEB 3 1981
RECEIVED

Sponsored jointly by

Alberta
ENVIRONMENT

Environment Canada
Environnement Canada

15th Floor, Oxbridge Place
9820 - 106 Street
Edmonton, Alberta, Canada
T5K 2J6

ALBERTA OIL SANDS ENVIRONMENTAL RESEARCH PROGRAM

RESEARCH REPORTS

These research reports describe the results of investigations funded under the Alberta Oil Sands Environmental Research Program. This program was designed to direct and co-ordinate research projects concerned with the environmental effects of development of the Athabasca Oil Sands in Alberta.

A list of research reports published to date is included at the end of this report.

Enquiries pertaining to the reports in the series should be directed to:

Research Management Division
15th Floor, Oxbridge Place
9820 - 106 Street
Edmonton, Alberta
T5K 2J6
(403) 427-3943

Differences in the Composition of Soils Under Open and Canopy Conditions at Two Sites Close-in to the Great Canadian Oil Sands Operation, Fort McMurray, Alberta

Project LS 3.4.1

AOSERP Report 97

This report may be cited as:

Baker, J. 1980. Differences in the composition of soils under open and canopy conditions at two sites close-in to the Great Canadian Oil Sands Operation, Fort McMurray, Alberta. Prep. for the Alberta Oil Sands Environmental Research Program by Canadian Forestry Service, Environment Canada. AOSERP Report 97. 24 pp.

The Hon. J.W. (Jack) Cookson
Minister of the Environment
222 Legislative Building
Edmonton, Alberta

and

The Hon. John Roberts
Minister of the Environment
Environment Canada
Ottawa, Ontario

Sirs:

Enclosed is the report "Differences in the Composition of Soils Under Open and Canopy Conditions at Two Sites Close-in to the Great Canadian Oil Sands Operation, Fort McMurray, Alberta".

This report was prepared for the Alberta Oil Sands Environmental Research Program, through its Land System under the Canada-Alberta Agreement of February 1975 (amended September 1977).

Respectfully,

W. Solodzuk, P.Eng.
Chairman, Steering Committee, AOSERP
Deputy Minister, Alberta Environment

A.H. Macpherson, Ph.D
Member, Steering Committee, AOSERP
Regional Director-General
Environment Canada
Western and Northern Region

DIFFERENCES IN THE COMPOSITION OF SOILS UNDER OPEN AND CANOPY CONDITIONS AT TWO SITES CLOSE-IN TO THE GREAT CANADIAN OIL SANDS OPERATION, FORT McMURRAY ALBERTA

DESCRIPTIVE SUMMARY

Atmospheric emissions from power plants, petroleum and gas plants, and other industrial activities may adversely affect the environment. Sulphur dioxide is a potentially hazardous atmospheric pollutant resulting from oil sands extraction. Although the present levels of emission do now show evidence of being hazardous to the environment (AOSERP Project LS 3.4 by Addison and Baker 1979), detailed studies undertaken in this project revealed that the site closer (2.3 km) to the Suncor plant showed degradation of ground lichens whereas the other site (5.3 km) did not show such degradation.

The soils sampled from these two sites were significantly different in several macronutrients. Also, the jackpine foliage from the closer site contained more sulphur than that from the distant site.

This report has been reviewed and accepted by the Alberta Oil Sands Environmental Research Program.

W.R. MacDonald, Ph.D
Director (1980-81)
Alberta Oil Sands Environmental
Research Program

101-06757/

DIFFERENCES IN THE COMPOSITION OF SOILS UNDER OPEN AND CANOPY CONDITIONS AT TWO SITES CLOSE-IN TO THE GREAT CANADIAN OIL SANDS OPERATION, FORT McMURRAY, ALBERTA

by

J. BAKER
Northern Forest Research Centre
Canadian Forestry Service

for

ALBERTA OIL SANDS ENVIRONMENTAL RESEARCH PROGRAM

Project LS 3.4.1

April 1980

TABLE OF CONTENTS

		Page
DECLARATION		ii
LETTER OF TRANSMITTAL		iii
DESCRIPTIVE SUMMARY		iv
LIST OF TABLES		viii
LIST OF FIGURES		ix
ABSTRACT		xi
ACKNOWLEDGEMENTS		xii
1.	INTRODUCTION	1
2.	MATERIALS AND METHODS	3
2.1	Soil Sampling	3
2.2	Foliar Sampling	5
3.	RESULTS AND DISCUSSION	6
3.1	Sulphur	6
3.2	Phosphorus	10
3.3	Soil Bases (Ca, Mg, and K)	13
3.4	Soil Sesquioxides (Al, Fe, and Ti)	13
4.	JACK PINE FOLIAGE ANALYSIS	15
5.	CONCLUSIONS	17
5.1	Soil Analysis	17
5.2	Foliar Analysis	17
6.	RECOMMENDATIONS	18
7.	REFERENCES CITED	19
8.	LIST OF AOSERP RESEARCH REPORTS	21

LIST OF TABLES

Page

1. Mean pH Values and Concentrations (ppm) in Soils from Horizons Under Jack Pine Cover and in Open Areas at Sites 1 and 2 Extracted by NH_4Cl 7

2. Mean Nutrient Concentrations (ppm) in Soils from Horizons Under Jack Pine Cover and in Open Areas at Sites 1 and 2 Extracted by 6N HCl 8

3. Mean Profile Concentrations, to the Next Whole Numbers (ppm) Under Jack Pine Cover and in Open Areas as Shown by NH_4Cl and 6N HCl Extraction 11

4. Mean Nutrient Status (ppm) in Current and One-Year-Old Jack Pine Growth (Needle Tissue) from Sites 1 and 2 16

LIST OF FIGURES

Page

1. Map of the AOSERP Study Area 2

2. Location of Gradient Sites in the Vicinity of the GCOS Operation 4

3. Wind Rose Direction Frequency for Mildred Lake 9

ABSTRACT

Soils sampled at Sites 1 and 2 [2.3 and 5.3 km south of the Great Canadian Oil Sands (GCOS) plant, Fort McMurray, respectively] were found to differ statistically with reference to several macronutrients. Soils under cover of a canopy and those in open areas differed considerably with respect to soil nutrients and pH. Differences in nutrient concentrations under open versus covered soils probably are due mainly to effects of the canopy; however, this may not fully explain soil sulphur and titanium concentrations.

Jack pine foliage sampled at Site 1 was more concentrated in S than that sampled at Site 2, suggesting the possibility that atmospheric sulphur may have played a role. Other foliar nutrient differences more likely are the result of other or natural phenomena.

To date, despite the higher S concentrations in both jack pine foliage and soils under the canopy at Site 1, there is no conclusive evidence that atmospheric pollutants emitted as a result of the overall exploration operation have as yet altered the existing soil nutrient regime of the study area, or are responsible for differences observed between the two sites.

ACKNOWLEDGEMENTS

The author wishes to extend his thanks to D.G. Caldwell and P. Addison [Northern Forest Research Centre (NFRC)] for their technical assistance throughout the investigation. Application is also directed to B. Chow (NFRC) and M. Kingsley [Canadian Wildlife Service (CWS)] for their assistance in the statistical treatment and computer processing of data.

This research project LS 3.4.1 was funded by the Alberta Oil Sands Environmental Research Program, a joint Alberta-Canada research program established to fund, direct, and co-ordinate environmental research in the Athabasca Oil Sands area of northeastern Alberta.

1. INTRODUCTION

Atmospheric emissions from power plants, petroleum and gas plants, and other industrial activities may adversely affect the environment. A potentially hazardous atmospheric pollutant resulting from the oil sands extraction is sulphur dioxide (SO_2). While as yet there appears to be little damage or danger to the environment from present levels of atmospheric SO_2, the fact that larger extraction plants are soon to be operational could alter markedly this situation.

This study was undertaken to increase the existing baseline information on a specific soil environment within the AOSERP study area (Figure 1) and its response to atmospheric pollutants arising from the oil sands exploration activities.

In view of the fact that Addison and Baker (1979) were unable to demonstrate any detrimental influence to the environment directly attributable to atmospheric pollutants, two sites due south of, and within 5.5 km of the Great Canadian Oil Sands (GCOS)[1] stack, on degraded dystic brunisol soils were studied in greater detail. The closer (2.3 km) of these sites showed some degradation of ground lichens whereas the other (5.3 km) did not. It was felt that, if pollutants arising from the GCOS operation were to demonstrate adverse effects, these would be observed most readily at these close-in sites where higher ambient SO_2 concentrations have been reported (Smith 1979).

Samples of both soils and jack pine foliage were taken and analyzed chemically and the results subjected to statistical analysis. This report records the results and possible explanations of these are offered.

---

[1]GCOS amalgamated with Sun Oil Company in August 1979, after the writing of this report was completed, to become Suncor, Inc.



Figure 1. Map of the AOSERP study area.

2. MATERIALS AND METHODS

Two sites (1 and 2, 2.3 and 5.3 km south of the GCOS plant, respectively) were selected and sampled with reference to soil and jack pine foliage. Both soils and vegetative cover were described previously by Addison (1976). Sites 1 and 2 represent Sites A and D, respectively, as described in Addison (1979) (Figure 2). Additional information regarding the soils and the vegetation and the positioning of sites within the study area was supplied by Addison and Baker (1979).

2.1 SOIL SAMPLING

Soil pits 2 x 1.5 x 1 m were dug immediately under a clump of pine trees (about four or five) and in an open space between clumps. The four walls of the pits (north, south, east, and west) were sampled at the litter surface (L-F-H), the surface of the Ae horizon (u-Ae), the lower boundary of the Ae (ℓ-Ae), the upper boundary of the Bm (u-Bm), the lower Bm boundary (ℓ-Bm), and in the mid-C horizon. The sampling depths within the profile corresponded to 6-0, 0-3, 12-15, 15-18, 33-36,and 61 cm at Site 1 and 3-0, 0-3, 7-10, 10-13, 39-42, and 61 cm at Site 2. These profile samples were analyzed separately.

Duplicate 5 and 10 g of field moist L-F-H and mineral soils were extracted for 4 h with 1N ammonia chloride (0.1 N in ethylene diaminetetraacetic acid) immediately followed by a 16 h extraction with 6N hydrochloric acid. All extractions were made by refluxing at 70°C under a partial vacuum of 50 kPa. In principle, the extraction procedure used was essentially that of Janssen (1958), who applied the method to the fractionation of soil nitrogen (N). His assumptions were that the salt extractable N and other soil nutrients were immediately or readily available for plant uptake and at least a portion of the acid extractable nutrients also would be available, arising as they do in part by the hydrolysis of organic matter and easily hydrolyzed mineral matter. However, the emphasis of the extraction technique in this study was not mainly in reference to availability but



Figure 2. Location of gradient sites in the vicinity of the GCOS operation.

rather was used as a means of demonstrating either natural or induced soil nutrient concentration differences associated with the sites under the several sampling regimes.

Analysis of the extracts included: sulphate-sulphur (SO_4-S) by the Dean modification of the Johnson-Nishita procedure as reported by Carson et al. (1972); phosphorus as the orthophosphate ($H_2PO_4^-$) by the molybdenum blue method (Jackson 1958); cationic species, calcium (Ca), magnesium (Mg), potassium (K), aluminum (Al), iron (Fe), and titanium (Ti), were evaluated by atomic spectrophotometry using an I.L. 251 unit and methods supplied by the manufacturer.

2.2 FOLIAR SAMPLING

Foliar samples were removed from five trees in close proximity to the soil pits at both sites. Current and one-year-old tissues were sampled in late fall from the upper three-whorl branches, washed with distilled water, dried at 75°C, and reduced to 60 mesh in a Wiley mill. Triplicate 0.10 to 0.15 g samples of prepared materials were ashed in a combustion flask which contained 25 mL of 0.5N HCl (0.18% in hydrogen peroxide) as an absorber. The flasks remained stoppered during and for 0.5 h after combustion to minimize loss. Aliquots of the absorbing solution were analyzed for SO_4-S, $H_2PO_4^-$, Ca, Mg, K, Al, Fe, and Mn using methods similar to those used in the soil analyses. Because of the combustion technique used, values recorded are total concentrations.

Since it is well known that vegetative materials can absorb gaseous atmospheric compounds directly through their leaves, the examination of jack pine foliage was included in a predominantly soils study to aid in the determination of the effects, if any, that atmospheric pollutants were having on the nutrient status, balance, and pH of soils in close proximity to the GCOS operation.

Statistical treatment of soil and foliar data was achieved by a 2^2 factorial analysis.

3. RESULTS AND DISCUSSION

3.1 SULPHUR

The factors considered in the experimental design, distance from GCOS stack, and cover, were statistically significant ($P < 0.01$) in the concentration and distribution of both soil S-forms in the profiles at the two sites. Since there was a strong interaction between the two factors ($P < 0.01$), it was apparent that they did not act independently of each other (Snedecor and Cochran 1956). However, data (Tables 1 and 2) indicated that the factors did not act similarly at the two sites. The NH_4Cl--extractable S concentration of the soil under cover at Site 1 was in considerable excess of that observed for the open area. The reverse held true for Site 2. Addison and Baker (1979), using the Huey Plate technique, reported that atmospheric SO_2 concentrations of Site 1 during the period of measurement were greater than those of Site 2. In addition, they reported that the various species checked for S all showed considerably higher concentrations at Site 1 than at Site 2, suggesting that Site 1 was the more heavily impinged. However, this increased atmospheric S concentration was not reflected in soil pH nor in the NH_4Cl--extractable soil S concentrations in samples from the open areas; the more distant Site 2 exhibited lower soil pH values and considerably higher S concentrations throughout the entire sampling depth (Table 1). In other words, despite supposedly greater atmospheric S concentrations, as suggested by the wind direction frequencies (Figure 3) which indicate a stronger impingement at Site 1, the open soils at Site 1 exhibited higher soil pH values and lower S concentrations than those at Site 2. However vegetation at Site 1 showed the higher S concentrations (Addison and Baker 1979). Since four separate profiles were sampled individually and analyzed, this anomaly hardly can be attributed to a sampling inconsistency. Consequently, the only reasonable explanation appeared to be that of a natural soil variation. Thus, the statistical significant difference in S concentration in the open soils of these sites may not exclusively be attributed directly to the GCOS operation.

Table 1. Mean pH values and nutrient concentrations † (ppm) in soils from horizons under jack pine cover and in open areas at Sites 1 and 2 (2.3 and 5.3 km south of the GCOS plant, respectively) extracted by NH_4Cl.[a]

Horizons	Cover	S	P	Ca	Mg	K	Al	Fe	Ti	Ph
Site 1										
L-F-H	Canopy	238.7	111.5	4334.9	495.5	643.2	689.3	1365.0	13.5	4.3
	Open	18.4	98.4	4675.2	541.9	529.6	547.0	1339.9	11.0	4.7
u-Ae	Canopy	9.0	12.0	83.7	10.5	16.8	45.6	340.8	5.8	4.4
	Open	0.6	23.0	149.3	19.1	27.1	122.8	330.2	6.8	4.9
ℓ-Ae	Canopy	9.7	13.9	72.5	9.5	17.6	56.8	392.0	7.0	4.4
	Open	0.3	31.8	130.4	17.6	32.8	151.7	537.2	8.1	4.9
u-Bm	Canopy	24.9	264.3	115.8	18.2	23.5	449.0	3489.9	39.5	4.4
	Open	1.7	23.2	102.8	16.9	36.1	642.6	1668.3	19.9	4.6
ℓ-Bm	Canopy	10.8	122.6	101.6	17.6	17.4	390.4	1299.1	6.6	4.7
	Open	0.1	23.2	102.5	14.3	18.1	206.9	327.5	4.0	4.5
C	Canopy	9.0	10.6	140.6	23.3	15.9	218.6	306.5	1.9	4.9
	Open	0.0	7.0	126.4	19.1	17.2	117.9	90.1	2.5	4.6
Site 2										
L-F-H	Canopy	48.3	57.3	2603.1	205.7	493.2	579.1	1002.5	9.3	4.0
	Open	125.4	76.9	3310.5	343.7	495.4	520.6	794.3	0.0	4.5
u-Ae	Canopy	0.0	6.9	80.6	8.9	14.5	56.4	255.6	2.9	4.1
	Open	22.2	2.6	104.6	9.3	25.5	60.9	51.1	0.0	3.9
ℓ-Ae	Canopy	0.0	8.5	64.4	7.7	15.3	68.9	284.1	7.4	4.0
	Open	21.7	5.3	92.5	10.8	21.2	77.1	87.7	5.0	4.2
u-Bm	Canopy	3.0	9.4	167.1	23.4	29.8	399.4	821.8	4.6	4.6
	Open	42.7	22.0	87.1	21.1	31.4	655.3	836.7	9.0	4.3
ℓ-Bm	Canopy	0.0	3.0	101.5	15.8	20.2	256.7	165.1	0.7	4.7
	Open	22.1	1.7	110.5	22.5	15.2	186.5	127.0	0.0	4.6
C	Canopy	0.0	2.1	76.9	19.8	10.0	45.2	157.1	0.0	4.8
	Open	18.9	3.5	33.0	8.2	6.8	65.0	180.3	1.0	4.4

[a] † each datum represents a mean of 8 replications.

Table 2. Mean nutrient concentrations † (ppm) of soils from horizons under jack pine cover and in open areas at Sites 1 and 2 (2.3 and 5.3 km south of the GCOS plant, respectively) extracted by 6N HCl.

Horizons	Cover	S	P	Ca	Mg	K	Al	Fe	Ti
Site 1									
L-F-H	Canopy	111.6	108.5	2098.2	191.1	184.2	749.7	2126.2	45.6
	Open	69.0	126.0	2253.4	257.6	204.7	1119.5	1882.1	68.6
u-Ae	Canopy	12.9	1.9	26.6	33.1	13.5	223.4	964.6	17.5
	Open	1.6	39.4	43.2	65.2	36.1	483.9	2164.9	22.7
ℓ-Ae	Canopy	12.8	3.6	27.5	36.3	14.3	269.4	1214.4	20.4
	Open	1.8	58.1	68.8	123.1	47.9	812.3	2728.4	31.1
u-Bm	Canopy	14.1	90.2	55.4	188.4	61.1	1251.4	4425.7	22.3
	Open	2.4	185.5	117.2	474.4	100.2	2655.8	6304.0	43.8
ℓ-Bm	Canopy	12.5	86.1	67.4	255.8	73.3	1697.3	6521.5	31.6
	Open	1.9	123.7	86.2	427.2	111.0	2515.4	7609.8	36.9
C	Canopy	12.6	80.8	64.8	284.3	97.6	1782.9	5922.3	32.1
	Open	1.0	64.6	83.6	380.6	106.5	1960.3	7060.1	39.8
Site 2									
L-F-H	Canopy	94.1	72.8	697.6	81.9	38.4	503.3	1306.2	30.5
	Open	58.6	113.6	902.1	133.0	117.6	695.4	989.7	42.4
u-Ae	Canopy	1.1	5.8	19.3	31.7	10.0	243.7	752.4	15.8
	Open	0.6	10.6	18.3	23.1	22.9	220.8	562.8	18.8
ℓ-Ae	Canopy	1.8	9.3	23.6	32.8	11.7	318.6	1230.5	21.9
	Open	0.6	23.6	35.7	57.8	34.6	433.6	1263.8	27.3
u-Bm	Canopy	7.4	201.9	71.9	335.8	80.9	2921.9	10393.1	50.9
	Open	2.8	178.2	81.7	498.7	102.0	4252.4	11010.6	53.6
ℓ-Bm	Canopy	2.8	51.5	53.9	210.8	98.6	2698.1	6503.3	33.5
	Open	2.0	57.4	52.3	390.1	106.0	2639.4	6216.4	34.6
C	Canopy	1.5	25.4	46.8	208.0	66.0	989.7	3513.4	21.1
	Open	0.5	22.8	38.6	193.3	57.1	690.0	1630.5	30.0

[a] † each datum represent a mean of 8 replications.



Figure 3. Wind rose of direction frequency for Mildred Lake (from Walmsley and Bagg 1977).

Sulphur concentrations in open soils detected by 6N HCl, although slightly greater at Site 1 and thus more in keeping with the reported atmospheric gradient, were for practical purposes similar. Of the S removed by both extractants from samples from open areas, a slightly greater proportion of that at Site 2 was sensitive to NH_4Cl. This tended to suggest that natural soil variation was responsible for the S concentration differences.

The soil S concentrations removed by both extractants from samples from under the canopy were consistently greater at Site 1 (Tables 1 and 2). This may be explained partly by the fact that atmospheric SO_2 concentrations were greater at Site 1. As a consequence, opportunity for the direct absorption of S may have been enhanced. Mudd (1975) reported the possibility of the entry of SO_2 through the stomata and its conversion to SO_4-S. Probably much of this S subsequently was deposited on the soil surface by stemflow, throughfall, litterfall, etc. (Curlin 1968; Mahendrappa 1974) and redistributed throughout the solum by natural nutrient cycling.

In this study, however, there may be another, perhaps more valid, explanation for the very high S concentrations observed under the canopy at Site 1. In the digging of the soil pit (and this pit only), lens of bitumen were encountered frequently in the C horizon. Consequently, much of, if not all, the elevated S concentrations noted for these soils could have been due to the presence of S-enriched bituminous compounds. Because of the lack of any consistent trend between sites, data (Table 3) tend to confirm the suggestion that soil S concentrations observed at both sites under the two sampling conditions were due mainly to natural soil differences.

3.2 PHOSPHORUS

Concentrations of soil phosphorus detected by the two extractants under both conditions of sampling were usually much greater in the soils of Site 1 (Table 1), the one exception being HCl--extractable P in the u-Bm horizon under the canopy (Table 2). While the statistical analysis indicated that P concentrations throughout the profiles

Table 3. Mean profile concentrations, to the nearest whole numbers (ppm), of nutrients under jack pine cover and in open areas as shown by NH_4Cl and 6N HCl extraction.

	NH_4Cl[a]		6N HCl[a]		Sum	
	Canopy	Open[b]	Canopy	Open[b]	Canopy	Open
Sulphur						
Site 1	51	4	30	13	81	17
Site 2	9	42	18	11	27	53
Calcium						
Site 1	808	881	390	442	1198	1323
Site 2	516	623	152	188	668	811
Magnesium						
Site 1	96	105	165	288	261	393
Site 2	47	69	167	216	214	285
Potassium						
Site 1	123	110	74	101	197	211
Site 2	97	99	51	74	148	173
Phosphorus						
Site 1	89	35	62	100	151	135
Site 2	15	19	62	68	77	87
Iron						
Site 1	1199	716	3529	4625	4728	5341
Site 2	448	346	3951	3612	4399	3958
Aluminum						
Site 1	308	315	996	1591	1304	1906
Site 2	234	261	1279	1489	1513	1750
Titanium						
Site 1	12	9	28	40	40	49
Site 2	4	3	29	35	33	38

[a] Extraction with NH_4Cl or 6N HCl.

[b] Type of cover.

of soils at Site 1, under both sampling regimes and for both extractants, were significantly different ($P < 0.01$) to those observed for the profiles of Site 2, it is hardly expected that this was due to P emissions from the GCOS stack. Addison and Baker (1979) have shown that such emissions are only minimal. A more likely explanation for these concentration differences is the proximity of Site 1 to the highway, which for many years was unpaved. Verry and Timmons (1977) reported a variety of cations and anions, including P, are carried into the atmosphere during soil disturbances either from natural events or man's activity. Consequently, much of the P detected at Site 1 may have come as a result of the deposition of dust during highway construction and use and from the removal of the surface mantle in the extraction of the oil sands. The use of salt on the highway during winter conditions also could be contributory.

An interesting feature of these data was seen in the effect of the canopy on P distribution within the solum. Concentrations of NH_4Cl--extractable P were smaller, although not significantly, in the L-F-H, u- and ℓ-Ae horizons under the canopy, whereas in the u- and ℓ-Bm they were significantly higher ($P < 0.01$:$P < 0.05$). This probably was the result of both increased vegetative uptake, especially from the upper horizons under the canopy, and increased downward movement as a result of increased water movement through these soils arising from foliar drip and stemflow which are normally more acid than incident rainfall (Baker et al. 1977). Data (Table 1) tend to substantiate this view.

The effect of canopy was even more pronounced on the HCl--extractable P concentrations. These were considerably lower in all horizons than comparable samples from open areas, except in the C. Significant differences were observed in the L-F-H and u- and ℓ-Ae ($P < 0.01$) (Table 2). The acid stemflow and throughfall entering the upper soil horizons over a period of time may have slowly solubilized and leached a portion of the soil P normally sensitive to HCl into deeper soil layers bringing about a new distribution within the solum.

3.3 SOIL BASES (Ca, Mg, and K)

With a few exceptions, concentrations of Ca, Mg, and K removed by NH_4Cl were greater in soils at Site 1. These cations usually were more concentrated in soils from open sampling areas than in soils under the canopy (Tables 1 and 2). As with soil P, the concentrations of these cations extractable by NH_4Cl were greater in the lower solum (u- and ℓ-Bm) than in the upper solum (u- and ℓ-Ae). This was not so noticeable in the profiles of the open areas. Probably this situation was brought about by a combination of vegetative uptake and increased leaching in soils under the canopy.

With reference to HCl--extractable bases, the status and distribution of these in the profile were similar to those extractable by NH_4Cl. However the concentrations were considerably higher throughout. Data (Table 1 and 2) reflected the fact that both distance from the GCOS operation and canopy were significant in the distribution and concentrations of soil bases within the soil profiles. These differences, while real, probably were the result of natural phenomena with little relationship to the oil sands extraction processes. Because of the soil surface disturbances in the oil sands area, soil particulates and dust deposition may have been largely responsible for these differences (Integ 1973; Verry and Timmons 1977).

3.4 SOIL SESQUIOXIDES (Al, Fe, and Ti)

With the exception of Ti, these cations are found in very high concentrations as data (Table 1 and 2) clearly show. Consequently, Al and Fe initially are of little value in the assessment of pollutant effects on the soil. If in time pollution is a real danger, especially that associated with SO_2, both of these cations but particularly Al (Hutchinson and Whitby 1976) become significant as they are excellent indicators of the acidic effects of pollution.

Titanium, on the other hand, is an immediately useful indicator of atmospheric pollution effects on soil and vegetation. The relatively low concentrations of both forms in soils make detection of Ti pollution effects more readily apparent.

While both NH_4Cl-- and HCl-- extractable Ti were found in greater concentrations in the profiles of Site 1, in view of the natural site differences already noted, it is impossible at this time to assume that these differences were related directly to pollutant effects of the GCOS operation.

4. JACK PINE FOLIAGE ANALYSIS

While three nutrients, S, Mg, and K, appeared in higher concentrations in current than in one-year growth (Table 4), only K ($P < 0.05$) was statistically significant. In conifer nutritional studies, Miller and Miller (1976) preferred current tissue sampled in late autumn from the topmost branch whorls of dominant and codominant trees. Stated advantages of this sampling location were that it may be precisely defined and resampled in subsequent years and that it was not greatly influenced by sociological factors. Raupach (1975), Lowry and Avard (1969), and Lowry (1968) were in agreement with this view.

Foliar concentrations of S and Mg were significantly different at the two sites ($P < 0.01$) with those at Site 1 being the greater. Foliar K concentrations at this site were significantly lower ($P < 0.05$) than those at Site 2. There did not appear to be any consistent relationship with soil concentrations except in the case of S. The increased S concentrations in tissues sampled at Site 1 may have been responsible for the increased Mg concentrations observed (Wallace 1961).

All other nutrients determined, P, Ca, Al, and Fe, were more concentrated in one-year-old tissues. Of these, Ca, Al, and Fe showed statistically significant differences ($P < 0.01$). Again no consistent trends with soil concentrations were noted. This suggested that factors in addition to soil nutrient concentrations were influential in governing nutrient concentrations in jack pine foliage.

Table 4. Mean † nutrient status (ppm) in current and one-year-old growth of jack pine (needle tissue) from Sites 1 and 2 (2.3 and 5.3 km south of the GCOS plant, respectively).[a]

Site	Age	S	P	Ca	Mg	K	Al	Fe	Mn
1	Current	1705.1	1194.6	2197.8	1920.4	4439.3	530.7	244.5	646.6
1	One-year	1570.4	1355.0	3430.5	1349.5	3700.3	699.5	355.8	881.7
2	Current	1246.0	1200.0	2279.7	1157.5	6136.2	592.2	218.1	535.6
2	One-year	1187.8	1116.1	3351.1	1075.2	4023.6	792.4	317.6	767.9

[a] † each datum represents a mean of three replications from each of five sample trees.

5. CONCLUSIONS

5.1 SOIL ANALYSIS

Detailed soil analysis which included NH_4Cl-- and HCl--extractable constituents indicated that, to date, atmospheric pollutants arising directly from the development of energy and from the oil sands recovery processes have made little impact on the soils of the study area. Statistically significant differences observed in the concentrations of various nutrients at the two sites appear to have been due mainly to natural variation. Surface mantle disturbances taking place in the oil sands area quite possibly contributed to these differences by soil and dust particulate deposition. Data obtained in this study are part of the baseline information to be used for comparative purposes in future studies.

5.2 FOLIAR ANALYSIS

Foliar concentrations of the various nutrients studied showed no consistent relationship with concentrations in the soil nor with distance from the GCOS plant; S, which is a known stack emission, may be an exception. Jack pine foliage samples from Site 1 exhibited greater S concentrations than those from Site 2. However, the actual cause or causes for this are as yet uncertain. Considerable variability in foliar nutrient concentrations between sample trees emphasized the need for more extensive sampling.

6. RECOMMENDATIONS

Site variability was large and further extensive work will be required before a reasonably accurate evaluation of pollution effects, if any, can be made. An increased sampling base and a more extensive statistical approach under the present field conditions of the study area, while useful, probably will not supply the necessary information required to make such an evaluation.

Soil and vegetative (jack pine) transplants could be introduced with profit into the oil sands area in a manner analogous to lichen transplants. These would have to be of known properties and characteristics and in sufficient numbers for good replication. They would have to be placed at strategic locations and in such a manner that artifact effects were minimal (i.e., effects produced by the methods used, etc.).

In view of the extensive surface mantle disturbances occurring in the oil sands area during the mining of the oil sands, wet and dry particulate gauges (nipher type) also should be employed in addition to the standard precipitation traps and Huey Plates. Particulate deposition arising from the mining and not the oil production processes may be a far more important consideration, especially with reference to soils and conifers.

Periodic examination of these introduced transplants probably would facilitate an early and fairly accurate appreciation of both the degree and advance of atmospheric pollution effects.

7. REFERENCES CITED

Addison, P. 1976. Soils of the air pollution benchmark sites in the Athabasca Oil Sands area. Can. For. Ser., Edmonton File Rep. 18 pp.

Addison, P., and J. Baker. 1979. Ecological benchmarking and biomonitoring for the detection of airborne pollutant effects on vegetation and soils, 1975 to 1978. Prep. for the Alberta Oil Sands Environmental Research Program by Northern Forest Research Centre, Canadian Forestry Service, Edmonton. Report 46. 51 pp.

Baker, J., D. Hocking, and M. Nyborg. 1977. Acidity of open and intercepted precipitation in forests and effects on forest soils in Alberta, Canada. Water, Air and Soil Pollution. D. Reidel Publishing Company. 7:449-460.

Carson, J.A., J.M. Crepin, and P. Nemunis Siugzdines. 1972. A sulfate-sulphur method used to delineate the sulfur status of soils. Can. J. Soil Sci. 52:278-281.

Curlin, J.W. 1968. Nutrient cycling as a factor in site productivity and forest fertilization, tree growth and forest soils. Pages 313-325 in Youngberg, C.T., and B.C. Davey (eds.), Tree growth and forest soils. Oregon State University Press, Corvallis, Oregon.

Hutchinson, T.C., and L.M. Whitby. 1976. Effects of acid rainfall on heavy metal particulates on a boreal forest ecosystem near the Sudbury smelting region of Canada. In Proc. First Intl. Sym. on acid precipitation and the forest ecosystem. For. Serv. U.S.D.A. Tech. Rep. NE 23. Upper Darby, Pa.

Intercontinental Engineering of Alberta Ltd. (Integ.). 1973. An environmental study of the Athabasca Tar Sands Alberta Environment. Edmonton, Alberta.

Jackson, M.L. 1958. Soil chemical analysis. Prentice-Hall Engelwood Cliffs, N.J.

Janssen, S.L. 1958. Tracer studies on nitrogen transformation in soil with special attention to mineralization - immobilization relationship. Annals of the Royal Agricultural College of Sweden. 24:105-361.

Lowry, G.L. 1968. Variations in nutrient content of black spruce needles due to crown position, needle age, crown class and season of sampling procedures. Pulp and Paper Research Institute of Canada, Pointe Claire, Quebec. 31 pp.

Lowry, G.L., and P.M. Avard. 1969. Nutrient content of black spruce and jack pine needles 111. Seasonal variations and recommended sampling procedures. Pulp and Paper Research Institute of Canada. Pointe Claire, Quebec. 54 pp.

Mahendrappa, M.K. 1974. Chemical composition of stemflow from some eastern Canadian tree species. Can. J. For. Res. 4:17.

Miller, H.G., and J.D. Miller. 1976. Analysis of needle fall as a means of assessing nitrogen status in pine. Forestry 49:57-61.

Mudd, J.B. 1975. Sulfur dioxide in response of plants to air pollution. Pages 9-22 in Mudd, J.B., and T.T. Kozlowski (eds.), Responses of plants to air pollution. Academic Press, N.Y.

Raupauch, M. 1975. Trace elements disorders in *Pinus* and their correction. Pages 353-367 in Nicholas, D.J.D. and E.R. Egan (eds.), Trace elements in soil-plant-animal systems. Academic Press. New York.

Smith, S.B. (ed.) 1979. Alberta Oil Sands Environmental Research Program Interim Report covering the period April 1975 to November 1978. Prep. by A.S. Mann, R.A. Hursey, R.T. Seidner, and B. Kasinska-Banas. Edmonton, Alberta. 101 pp.

Snedecor, G.W., and W.G. Cochran. 1956. Statistical methods. The Iowa State College Press, Ames, Iowa.

Verry, E.S., and D.R. Timmons. 1977. Precipitation nutrients in the open and under two forests in Minnesota, U.S.A. Can. J. For. Res. 7:112-119.

Wallace, T. 1961. The diagnosis of mineral deficiencies in plants. Her Majesty's Stationery Office, London. 12 pp.

Walmsley, J.L., and D.L. Bagg. 1977. Calculations of annual averaged sulphur dicxide concentrations at ground level in the AOSERP study area. Prep. for the Alberta Oil Sands Environmental Research Program by Fisheries and Environment Canada, Atmospheric Environment Service. AOSERP Report 19. 40 pp.

8. AOSERP RESEARCH REPORTS

1.		AOSERP First Annual Report, 1975
2.	AF 4.1.1	Walleye and Goldeye Fisheries Investigations in the Peace-Athabasca Delta--1975
3.	HE 1.1.1	Structure of a Traditional Baseline Data System
4.	VE 2.2	A Preliminary Vegetation Survey of the Alberta Oil Sands Environmental Research Program Study Area
5.	HY 3.1	The Evaluation of Wastewaters from an Oil Sand Extraction Plant
6.		Housing for the North--The Stackwall System
7.	AF 3.1.1	A Synopsis of the Physical and Biological Limnology and Fisheries Programs within the Alberta Oil Sands Area
8.	AF 1.2.1	The Impact of Saline Waters upon Freshwater Biota (A Literature Review and Bibliography)
9.	ME 3.3	Preliminary Investigations into the Magnitude of Fog Occurrence and Associated Problems in the Oil Sands Area
10.	HE 2.1	Development of a Research Design Related to Archaeological Studies in the Athabasca Oil Sands Area
11.	AF 2.2.1	Life Cycles of Some Common Aquatic Insects of the Athabasca River, Alberta
12.	ME 1.7	Very High Resolution Meteorological Satellite Study of Oil Sands Weather: "A Feasibility Study"
13.	ME 2.3.1	Plume Dispersion Measurements from an Oil Sands Extraction Plant, March 1976
14.		
15.	ME 3.4	A Climatology of Low Level Air Trajectories in the Alberta Oil Sands Area
16.	ME 1.6	The Feasibility of a Weather Radar near Fort McMurray, Alberta
17.	AF 2.1.1	A Survey of Baseline Levels of Contaminants in Aquatic Biota of the AOSERP Study Area
18.	HY 1.1	Interim Compilation of Stream Gauging Data to December 1976 for the Alberta Oil Sands Environmental Research Program
19.	ME 4.1	Calculations of Annual Averaged Sulphur Dioxide Concentrations at Ground Level in the AOSERP Study Area
20.	HY 3.1.1	Characterization of Organic Constituents in Waters and Wastewaters of the Athabasca Oil Sands Mining Area
21.		AOSERP Second Annual Report, 1976-77
22.		Alberta Oil Sands Environmental Research Program Interim Report to 1978 covering the period April 1975 to November 1978
23.	AF 1.1.2	Acute Lethality of Mine Depressurization Water on Trout Perch and Rainbow Trout
24.	ME 1.5.2	Air System Winter Field Study in the AOSERP Study Area, February 1977.
25.	ME 3.5.1	Review of Pollutant Transformation Processes Relevant to the Alberta Oil Sands Area

26.	AF 4.5.1	Interim Report on an Intensive Study of the Fish Fauna of the Muskeg River Watershed of Northeastern Alberta
27.	ME 1.5.1	Meteorology and Air Quality Winter Field Study in the AOSERP Study Area, March 1976
28.	VE 2.1	Interim Report on a Soils Inventory in the Athabasca Oil Sands Area
29.	ME 2.2	An Inventory System for Atmospheric Emissions in the AOSERP Study Area
30.	ME 2.1	Ambient Air Quality in the AOSERP Study Area, 1977
31.	VE 2.3	Ecological Habitat Mapping of the AOSERP Study Area: Phase I
32.		AOSERP Third Annual Report, 1977-78
33.	TF 1.2	Relationships Between Habitats, Forages, and Carrying Capacity of Moose Range in northern Alberta. Part I: Moose Preferences for Habitat Strata and Forages.
34.	HY 2.4	Heavy Metals in Bottom Sediments of the Mainstem Athabasca River System in the AOSERP Study Area
35.	AF 4.9.1	The Effects of Sedimentation on the Aquatic Biota
36.	AF 4.8.1	Fall Fisheries Investigations in the Athabasca and Clearwater Rivers Upstream of Fort McMurray: Volume I
37.	HE 2.2.2	Community Studies: Fort McMurray, Anzac, Fort MacKay
38.	VE 7.1.1	Techniques for the Control of Small Mammals: A Review
39.	ME 1.0	The Climatology of the Alberta Oil Sands Environmental Research Program Study Area
40.	WS 3.3	Mixing Characteristics of the Athabasca River below Fort McMurray - Winter Conditions
41.	AF 3.5.1	Acute and Chronic Toxicity of Vanadium to Fish
42.	TF 1.1.4	Analysis of Fur Production Records for Registered Traplines in the AOSERP Study Area, 1970-75
43.	TF 6.1	A Socioeconomic Evaluation of the Recreational Fish and Wildlife Resources in Alberta, with Particular Reference to the AOSERP Study Area. Volume I: Summary and Conclusions
44.	VE 3.1	Interim Report on Symptomology and Threshold Levels of Air Pollutant Injury to Vegetation, 1975 to 1978
45.	VE 3.3	Interim Report on Physiology and Mechanisms of Air-Borne Pollutant Injury to Vegetation, 1975 to 1978
46.	VE 3.4	Interim Report on Ecological Benchmarking and Biomonitoring for Detection of Air-Borne Pollutant Effects on Vegetation and Soils, 1975 to 1978.
47.	TF 1.1.1	A Visibility Bias Model for Aerial Surveys for Moose on the AOSERP Study Area
48.	HG 1.1	Interim Report on a Hydrogeological Investigation of the Muskeg River Basin, Alberta
49.	WS 1.3.3	The Ecology of Macrobenthic Invertebrate Communities in Hartley Creek, Northeastern Alberta
50.	ME 3.6	Literature Review on Pollution Deposition Processes
51.	HY 1.3	Interim Compilation of 1976 Suspended Sediment Date in the AOSERP Study Area
52.	ME 2.3.2	Plume Dispersion Measurements from an Oil Sands Extraction Plan, June 1977

53.	HY 3.1.2	Baseline States of Organic Constituents in the Athabasca River System Upstream of Fort McMurray
54.	WS 2.3	A Preliminary Study of Chemical and Microbial Characteristics of the Athabasca River in the Athabasca Oil Sands Area of Northeastern Alberta
55.	HY 2.6	Microbial Populations in the Athabasca River
56.	AF 3.2.1	The Acute Toxicity of Saline Groundwater and of Vanadium to Fish and Aquatic Invertebrates
57.	LS 2.3.1	Ecological Habitat Mapping of the AOSERP Study Area (Supplement): Phase I
58.	AF 2.0.2	Interim Report on Ecological Studies on the Lower Trophic Levels of Muskeg Rivers Within the Alberta Oil Sands Environmental Research Program Study Area
59.	TF 3.1	Semi-Aquatic Mammals: Annotated Bibliography
60.	WS 1.1.1	Synthesis of Surface Water Hydrology
61.	AF 4.5.2	An Intensive Study of the Fish Fauna of the Steepbank River Watershed of Northeastern Alberta
62.	TF 5.1	Amphibians and Reptiles in the AOSERP Study Area
63.	ME 3.8.3	Analysis of AOSERP Plume Sigma Data
64.	LS 21.6.1	A Review and Assessment of the Baseline Data Relevant to the Impacts of Oil Sands Development on Large Mammals in the AOSERP Study Area
65.	LS 21.6.2	A Review and Assessment of the Baseline Data Relevant to the Impacts of Oil Sands Development on Black Bears in the AOSERP Study Area
66.	AS 4.3.2	An Assessment of the Models LIRAQ and ADPIC for Application to the Athabasca Oil Sands Area
67.	WS 1.3.2	Aquatic Biological Investigations of the Muskeg River Watershed
68.	AS 1.5.3 AS 3.5.2	Air System Summer Field Study in the AOSERP Study Area, June 1977
69.	HS 40.1	Native Employment Patterns in Alberta's Athabasca Oil Sands Region
70.	LS 28.1.2	An Interim Report on the Insectivorous Animals in the AOSERP Study Area
71.	HY 2.2	Lake Acidification Potential in the Alberta Oil Sands Environmental Research Program Study Area
72.	LS 7.1.2	The Ecology of Five Major Species of Small Mammals in the AOSERP Study Area: A Review
73.	LS 23.2	Distribution, Abundance and Habitat Associations of Beavers, Muskrats, Mink and River Otters in the AOSERP Study Area, Northeastern Alberta
74.	AS 4.5	Air Quality Modelling and User Needs
75.	WS 1.3.4	Interim Report on a Comparative Study of Benthic Algal Primary Productivity in the AOSERP Study Area
76.	AF 4.5.1	An Intensive Study of the Fish Fauna of the Muskeg River Watershed of Northeastern Alberta
77.	HS 20.1	Overview of Local Economic Development in the Athabasca Oil Sands Region Since 1961.
78.	LS 22.1.1	Habitat Relationships and Management of Terrestrial Birds in Northeastern Alberta

79.	AF 3.6.1	The Multiple Toxicity of Vanadium, Nickel, and Phenol to Fish.
80.	HS 10.2 & HS 10.1	History of the Athabasca Oil Sands Region, 1980 to 1960's. Volumes I and II.
81.	LS 22.1.2	Species Distribution and Habitat Relationships of Waterfowl in Northeastern Alberta.
82.	LS 22.2	Breeding Distribution and Behaviour of the White Pelican in the Athabasca Oil Sands Area.
83.	LS 22.2	The Distribution, Foraging Behaviour, and Allied Activities of the White Pelican in the Athabasca Oil Sands Area.
84.	WS 1.6.1	Investigations of the Spring Spawning Fish Populations in the Athabasca and Clearwater Rivers Upstream from Fort McMurray; Volume I.
85.	HY 2.5	An intensive Surface Water Quality Study of the Muskeg River Watershed. Volume I: Water Chemistry.
86.	AS 3.7	An Observational Study of Fog in the AOSERP Study Area.
87.	WS 2.2	Hydrogeological Investigation of Muskeg River Basin, Alberta
88.	AF 2.0.1	Ecological Studies of the Aquatic Invertebrates of the Alberta Oil Sands Environmental Research Program Study Area of Northeastern Alberta
89.	AF 4.3.2	Fishery Resources of the Athabasca River Downstream of Fort McMurray, Alberta. Volume I
90.	AS 3.2	A Wintertime Investigation of the Deposition of Pollutants around an Isolated Power Plant in Northern Alberta
91.	LS 5.2	Characterization of Stored Peat in the Alberta Oil Sands Area

These reports are not available upon request. For further information about availability and location of depositories, please contact:

Alberta Oil Sands Environmental Research Program
15th Floor, Oxbridge Place
9820 - 106 Street
Edmonton, Alberta
T5K 2J6